



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Hylocichla ustulata swainsoni. OLIVE-BACKED THRUSH.—May 19, 1920, among the hills in Breathitt Co.

Planesticus migratorius migratorius. ROBIN.—A common bird May 8 to 23 and September 23 to 26, in the level counties but three only seen about the town of Jackson, Breathitt Co., May 19, 1920, and not found elsewhere in the hills

Sialia sialis sialis. BLUEBIRD.—Several noted in about all counties visited.

Highland Park, Reservoir Ave., Rochester, N. Y.

A NEW BURROWING OWL FROM COLOMBIA.

BY WITMER STONE.

IN a collection of Colombian birds recently received by the Academy of Natural Sciences of Philadelphia from Mr. M. A. Carriker, Jr., are two specimens of a Burrowing Owl which appear to represent an undescribed form.

They are nearest geographically to *Speotyto cunicularia tolimae* described by the writer¹ from the plains of Tolima, but curiously enough while that form is the darkest Burrowing Owl that I have seen from South America the present is among the lightest. It may be described as follows:

***Speotyto cunicularia carrikeri* subsp. nov.**

Type. No. 73700 Coll. Acad. Nat. Sci., Philadelphia. Palmar, Boyaca, Colombia. April 24, 1917, coll. M. A. Carriker, Jr.

Description. Resembles almost exactly in general coloration specimens of *S. c. hypogaea*, from California, and similar to the large pale birds of southern South America, but it is of about the same dimensions as the small dark colored *S. c. tolimae* of Colombia.

The outermost tail feathers are white with remains of four dusky bars mainly on the outer web and reduced to a mere streak in the case of the uppermost; on the next three pairs the bars are broad and well defined on the outer web, but the inner web is largely buffy white, the distal bars sometimes partly developed. The outer feathers are equally barred on both webs.

This bird is much worn and the light spots above and on the wings are white.

The other example (No. 73699) is marked as a female and is distinctly tawny, the tail more heavily barred and all light spots on tail, wings and back, strongly suffused with tawny. There is, however, in neither specimen a suggestion of the almost chocolate brown of *S. c. tolimae*.

The type measures as follows: Wing 173 mm, tail 78, tarsus 46. exposed culmen (from cere) 14.

¹ Stone, Proc. Acad. Nat. Sci., Phila., 1899. p. 304.